

Aging and Health Status of Elderly in Latin America and the Caribbean: Preliminary Findings

Alberto Palloni · Mary McEniry

Published online: 5 October 2006

© Springer Science + Business Media, Inc. 2006

Abstract Aging in Latin America and the Caribbean will not proceed along known paths already followed by more developed countries. In particular, the health profile of the future elderly population is less predictable due to factors associated with their demographic past that may haunt them for a long time and make them more vulnerable, even if economic and institutional conditions turn out to be better than what they are likely to be. This paper answers a set of questions regarding the nature and determinants of health status among the elderly in Latin America and the Caribbean using SABE (Survey on Health and Well-Being of Elders), a cross-sectional representative sample of over 10,000 elderly aged 60 and above in private homes in seven major cities in Latin America and the Caribbean. We examine health outcomes such as self-reported health, functional limitations—Activities of Daily Living (ADL's) and Instrumental Activities of Daily Living (IADL's), obesity (ratio of weight in kilograms to the square of height in centimeters), and self-reported chronic conditions (including diabetes). The findings include: (a) Countries differ in self-reported health but exhibit much less differences in terms of functional limitations. The number of chronic conditions increase with age and is higher among females than among males; (b) On average SABE countries display levels of self-reported diabetes (and obesity) that are as high if not higher than those found in the US; (c) There is evidence, albeit weaker than expected, suggesting deteriorated health and functional status in the region; (d) There is important evidence pointing toward rather strong inequalities (by education and income) in selected health outcomes. Preliminary findings from SABE confirm that Latin America and the Caribbean display peculiarities in the health profile of elderly, particularly with regard to diabetes and obesity. It is important that new policy initiatives begin to seriously target

A. Palloni

Center for Demography and Ecology, University of Wisconsin-Madison, Madison, WI, USA

M. McEniry

Center for Demography of Health and Aging (CDHA), University of Wisconsin-Madison, Madison, WI, USA

A. Palloni (✉)

Room 4434 Social Science Building, Madison, WI 53706, USA

e-mail: palloni@ssc.wisc.edu

the region's elderly, especially with an emphasis on the prevention and treatment of diabetes and obesity.

Keywords Aging · Diabetes · Health outcomes · Latin America

Introduction

We use a new and rich data set on elderly people in seven countries of Latin America and the Caribbean to document salient properties of the health status profile of elderly in the region and to answer a set of three questions regarding its nature and determinants. We use this new data set to explore conditions of elderly in the various subpopulations, a task made impossible up to now due to lack of information. Although we start out with a conjecture about the health status of elderly in the region and we use it to guide us in the analysis of information, our most important goal is not one of theory-building but of rich description to identify the main areas of interest that warrant further research.

Health status of elderly is but one among a handful of characteristics that make the aging process of the region a distinct one. There are four features that set the aging processes in the region of Latin America and the Caribbean apart from others. These features are singularities in an otherwise standard landscape of aging. They are potentially important for, even if verified only partially, they may generate a unique mixture of problems requiring vastly different policies from those undertaken elsewhere. What follows is a succinct review of these features.¹

Speed of Aging

The speed of demographic aging in Latin America and the Caribbean will be unprecedented. The time it will take a typical country in Latin America and the Caribbean to attain a substantial fraction of people above age 60, say around 15%, from current levels of around 8% is less than two fifths the length of time it took the US, and between one fifth and two fifths of the time it took an average Western European country to attain similar levels (Palloni, Pinto, & Pelaez, 2002; Kinsella & Velkoff, 2001). By 2030, in many countries in Latin America and the Caribbean, the number ages 60+ will be 2-1/2 to 3-1/2 times as large as it was in 2000. Barring unexpected demographic upheavals we should expect that for the next three to five decades the speed of aging in the region will continue on a singularly fast course, a result of the momentum of demographic force set in motion long ago.

Dislocation Between Aging and Standards of Living

Fast and demographically premature aging takes place in countries that have not had the time, the fortune, or wherewithal to generate sustained high standards of living. Comparisons between the wealthiest countries in Latin America, on the one hand, and the US, Sweden or Japan, on the other, are revealing. First, even optimistic projections of growth in GNP per capita imply that when the fraction of elderly people begins to exceed 10%, countries of the region will attain no more than a small fraction (one tenth or thereabouts) of the levels of GNP per capita enjoyed by developed countries when they were reaching similar levels of aging.

¹ A more thorough examination of the aforementioned features can be found in Palloni et al., 2002.

Second, adopting an admittedly optimistic forecast with a fairly agile process of economic growth, driven by annual rates of increase in GNP per capita of about 0.030 (about 15% higher than the average in the region during the last 50 years!) does not help much. Indeed, even in this rosy scenario, no appreciable fraction of time during which the countries of the region are aging rapidly will be characterized by a GNP per capita exceeding \$10,000, an admittedly modest amount. By comparison, Sweden spent about 77% of the time during which rapid aging was occurring enjoying higher standards of living; the US spent 95% of the time, and Japan 100% of the time. Barring unprecedented economic conjunctures, the fate of countries in Latin America and the Caribbean will be dominated by rapid aging paired with precarious standards of living. The comparisons are, of course, even more disheartening had we chosen as reference not the wealthiest but the poorest countries in the region.

Socio-Political Context and Aging: Institutional Volatility

An even more startling and generalized reality in the Latin American and Caribbean region has to do with the relation between speed and magnitude of aging, on the one hand, and the social and political contexts within which the process is taking place, on the other. First, a traditional order whereby elderly well-being rests on the shoulders of the younger generation is being gradually subverted by shifts in norms regulating living arrangements and by rapid fertility declines (Devos, 1990; Devos & Palloni, 2002; Palloni, 2001). Admittedly, traditional living arrangements crumbled in North America and Western Europe as well, but the phenomenon occurred well before the onslaught of rapid aging (Palloni, 2001; Ruggles, 1996). In countries in the Latin American and Caribbean region the safety net articulated around families and kin relations is being dismantled *concurrently* with rapid aging. This leaves little room for error and no time to seek adequate substitutes.

Second, aging is occurring in a fragile institutional environment, one where the bulk of sources guaranteeing minimum levels of social and economic support for the elderly are being reformulated, reformed, and in most cases, eliminated. A good example of this is the sustained and widespread drive toward reform of social security systems (Mesa-Lago, 1994; Barrientos, 1997; Klinsberg, 2000). In all cases the reforms are designed to replace pay-as-you go systems that operated uninterruptedly in many of these countries since World War I, with privatization schemes. New plans will supplant a system which, though flawed, was successful in reducing inequalities and protecting the most vulnerable segments of the elderly population. Income receipts of those retiring from the labor force during the first 10 years of the XXIst century will depend on clauses to stop-gap a prolonged transition to the instauration of the new system. An important fraction of these cohorts, but especially older women, received minimal earnings throughout their occupational careers and could not possibly accumulate sufficient wealth to secure safe standards of living. The combination of sheer growth of the elderly population and even an expected burden of disease among the elderly (see below) will result in an increase in the demand for health services precisely during a time when access to health care shrinks and becomes more expensive under the onslaught of privatization schemes.

In summary: no country in the Latin American and Caribbean region is blessed with institutional contexts designed to cope with changed demands from a growing elderly population. In almost all cases a highly compressed aging process will take place in the midst of weak economic performance, tense intergenerational relations, fragile institutional contexts, and shrinking access to medical and health care services.

Health Status

Birth cohorts who reach age 60 and above after 1990 in this region are unique in that they are largely composed by individuals who survived as a result of medical interventions in early childhood and largely in the absence of significant improvements in standards of living. It is estimated that between 50 to 70% of the mortality decline that took place after 1945 was associated with medical interventions (Preston, 1976; Palloni & Wyrick, 1981). The remaining decline was probably associated with better standards of living, increased knowledge about exposure and resistance to illnesses, and assorted other factors. Furthermore, a large fraction of these gains were concentrated early in the life of individuals, between birth and age 5 or 10.

A potential consequence of this pattern of mortality decline is that average levels of frailty among members of cohorts blessed by new medical technology could be higher (and so will its variance) than if mortality had remained constant or had declined due to improved standards of living. This is so for two reasons. *First*, the lives saved by the mortality decline were certainly not random relative to conditions affecting health status (Vaupel, Manton, & Stallard, 1979). Indeed, they are more likely to have been drawn from populations exposed to higher risks, those whose morbidity and mortality experiences were dominated by exposure to infectious diseases and lack of adequate early nutrition.

Second, since most childhood morbid conditions responsible for higher mortality before the interventions continued to affect children, albeit with reduced lethality, their influence and aftermath must have been felt by a growing number of survivors, all drawn from high mortality subpopulations. This has important implications *if early childhood conditions exert an impact on adult health and mortality*. Although evidence that early childhood conditions affect adult health is far from water-tight, it is clearly mounting fast and cannot be ignored. The literature on the subject focuses on *three types of factors* that have frequently been examined in this literature. This literature is split in several strands and we attempt to do justice to each of them (Elo & Preston, 1992; Schaffer, 2000; Barker, 1998; Kuh & Ben-Shlomo, 2004; Hertzman, 1994; Cynader, 1994). The first is most closely associated with the work of Barker and concentrates on the sequelae of processes that may start in utero or develop shortly before and/or around birth (“fetal origin hypothesis”). In general, these effects develop as a result of either fixed traits that individuals are born with or, most interestingly, of stresses and uneven development of physiological systems that follow periods of moderate and severe deprivation and that remain latent until late in life. Thus, unless one has markers of early deprivation, there is little that can be done to falsify the conjecture. Some of these markers have already been used (birth weight, placental weight, length of gestation, length of recently born etc...). None of these markers is available to us in the data we will use in this paper. Instead we rely on indirect measures of early nutritional status, including height (adjusted for age), knee height (a proxy for leg length), and the ratio of hip to waist circumference. These measures have already been used with some success by nutritionists as surrogates measures among adults. A second strand in the literature identifies episodes of illnesses in early childhood as the cornerstone responsible for the late onset of some chronic diseases. The best known example of this is the relation between rheumatic heart fever—a common infectious disease in developing countries at least prior to the massive mortality decline that took place after World War II,—and the onset of heart disease. Because all the data from SABE contain information on retrospectively recalled childhood diseases, we can at least attempt to assess the size of the effects. The strategy is by no means optimal since not only we must deal with faulty recall but

there is a serious selection problem we cannot address (for example, individuals with the most serious cases of rheumatic fever may not have survived to be in our sample). A third strand is more general and attempts to find broad associations between socioeconomic conditions experienced in early childhood and adult health status. This type of work is a roundabout way to find some of the connections identified by the other two strands. Thus, finding an association between SES early in life and health status among the elderly may simply reflect the relation between current or recent SES and health. For the most part this kind of work aims at finding *net effects of early SES on adult or health, that is, those that remain after appropriately controlling for current or recent SES*. The interpretation of the net effects conventionally invokes either the existence of Barker-effects or the influence of early illnesses. All the data sources we will use enable us to do tests for this as they contain retrospective evaluation of markers of early childhood poverty, deprivation and SES.

If any of these mechanisms turns out to have more than modest effects, increases in frailty among elderly whose earlier experiences fits the description provided above, are likely to be pronounced.² This means that the health status composition of elderly in Latin America and the Caribbean in general, should be worse relative to what would have been had the growth of the more recent and forthcoming cohorts of elderly been associated, as was the case in more developed countries, with improving standards of living. Our understanding of the relations between early childhood exposures and adult health status is still too primitive to enable us to establish precise predictions or conjectures regarding the nature of expected health impairments. But this conjecture can at least be used as a guiding torch to explore the evidence available to us.

Finally, it is neither a mystery nor a novelty that the regimes of morbidity and mortality experienced by elderly people in developing countries are unusual. First, as one would expect (Omran, 1982) there is an expansion of chronic conditions, such as heart and lung disease, cancers, diabetes, and arthritis, and elderly people continue to be assaulted by significant levels of infectious diseases (Frenk, Frejka, Bobadilla, Stern, & Sepulveda, 1991). We simply do not know what the health effects of exposure to highly interactive environments like these are. What should one expect, for example, under conditions where elderly people are simultaneously weakened by malaria and exposed to higher risks of congestive heart disease? Or, where increases in diabetes due to the adoption of a westernized diet (Popkin, 1993; Albala, Kain, Burrows, & Diaz, 2000) are combined with recurrent intestinal infections and high prevalence of respiratory TB? What are the implications of a mixed mode of exposure for comorbidities, disability and impairments among the elderly? What are the implications for treatment? What effects does it have on demands for health care?

The strength of evidence supporting the existence of the aforementioned features of the aging process in the Latin American and Caribbean region is heterogeneous. There is considerably more data sustaining hypotheses about the first and second features than there is for the third but especially for the fourth. It is precisely this fourth feature that constitutes the focus of this paper. We do this by turning our focus to self-reported health, functional limitations and chronic conditions, especially diabetes.

Up to now very little was known about adult health in the region and, therefore, virtually no conjecture or hypothesis could be thoroughly investigated. We can now use a unique, newly released data set on elderly people living in seven major cities in countries of Latin

² The argument holds, of course, if we assume that the effects of mortality selection are lonely mild and if the effects of changes in behavioral profiles and medical technology (exogenous or not) are only weak.

America and the Caribbean to answer the following three questions: (a) what is the characteristic health profile of the elderly in the region? (b) How does this profile compare with other known profiles, such as the one in the US? (c) Is there any evidence of deteriorated health and functional status, as expected if some of the conjectures proposed before are on the mark?

Subjects and Methods

SABE

SABE (SABE, 2003) is a data collection project anchored in seven major cities (six of them capital cities) of the region: Buenos Aires (Argentina), Bridgetown (Barbados), San Paulo (Brazil), Santiago (Chile), Havana (Cuba), Mexico City (Mexico) and Montevideo (Uruguay). All seven surveys were administered to representative samples of populations aged 60 and above in each city and were strictly comparable though translated to three different languages (Spanish, Portuguese and English). In some cases, interviewers selected a target older person and his/her surviving spouse. All sample frames were drawn either from recent population censuses or from nationally representative surveys carried out periodically in the capital cities of the region.³ The fieldwork took place between June 1999 and June 2000 and a preliminary final report was completed in December of 2002. An important feature of the survey is that, with two exceptions (Buenos Aires and Montevideo), the rates of response were significantly higher than those in similar surveys in other countries. Table 1 displays basic information on sample sizes, rates of response, as well selected dimensions of the demographic profile (composition by age, sex, marital status, race) and of the socioeconomic composition of the samples (by education). As shown elsewhere the basic demographic profile accords well with national figures.^{4,5} Table 2 displays information on a few health-related characteristics that will be the object of study in this paper, namely, self-reported health status, Activities of Daily Living (ADL), Instrumental Activities of Daily Living (IADL), chronic conditions, and anthropometric measures.⁶

HRS

The University of Michigan Health and Retirement Study (HRS) surveys more than 22,000 Americans over the age of 50 every 2 years. The study paints an emerging portrait of an aging America's physical and mental health, insurance coverage, financial status, family

³ For more information on the nature of the samples see Palloni and Pelaez, 2002 and Albala et al., 2005.

⁴ Because all samples are urban samples, our ability to generalize to the total population is impaired. However, readers should bear in mind that the proportion of the total population living in urban areas in these countries is substantial, varying from close to 100% in Barbados to about 74 or 75% in Mexico and Cuba, respectively (United Nations, 2000). This suggests that our results should not be too different from what we would have obtained had SABE been based on national samples. And, indeed, it has been shown that the demographic profile at least of the samples is quite close to national averages (Palloni & Pelaez, 2002).

⁵ In the rest of the paper we refer use the words "country" or "city" to refer to the *city samples*. By using the word country we are in no way assuming that the SABE data are exactly representative of elderly populations in each of the countries who participated in the project.

⁶ The definition of ADL, IADL, and self-reported conditions selected for study in this paper appears in the Appendix. They are strictly comparable to those used in other surveys of elderly, particularly the Health and Retirement Survey (HRS) (2002).

Table 1 Basic sample information.

Condition/ variable	Overall (<i>n</i> = 10,902)	Argentina (<i>n</i> = 1,043)	Barbados (<i>n</i> = 1,808)	Brazil (<i>n</i> = 2,143)	Chile (<i>n</i> = 1,306)	Cuba (<i>n</i> = 1,905)	Mexico (<i>n</i> = 1,247)	Uruguay (<i>n</i> = 1,450)
Response rate		0.60	0.85	0.85	0.84	0.95	0.85	0.66
Age	72 (8)	71 (7)	72 (8)	73 (8)	72 (8)	72 (9)	70 (8)	71 (7)
60–64	23%	23%	19%	20%	22%	25%	31%	22%
65–69	23%	24%	23%	18%	25%	21%	25%	25%
70–74	19%	24%	21%	16%	19%	18%	18%	23%
75–79	17%	15%	17%	22%	16%	13%	13%	17%
80–84	11%	8%	11%	14%	10%	11%	8%	9%
85+	8%	5%	9%	10%	8%	11%	6%	5%
Gender								
Females	62%	63%	60%	59%	66%	63%	59%	63%
Education								
Primary	71%	71%	77%	85%	68%	57%	74%	65%
Secondary	20%	23%	18%	5%	24%	37%	11%	21%
Higher	9%	6%	5%	10%	9%	7%	15%	14%
Race								
White	55%	–	5%	71%	43%	63%	–	90%
Black	34%	–	93%	16%	1%	36%	–	4%
Mestizo	6%	–	0.84%	8%	30%	–	–	6%
Other	4%	–	1%	5%	26%	0.58%	–	0.07%
Marital status								
Never married	7%	6%	18%	5%	7%	3%	4%	4%
Married/union	46%	43%	45%	52%	44%	37%	54%	49%
Separated	9%	9%	9%	6%	13%	13%	9%	6%
Widowed	34%	42%	24%	35%	36%	35%	32%	37%
Divorced	4%	0.96%	5%	1%	0.39%	11%	1%	5%

Source: SABE data, respondents ages 60 and above. Numbers in parentheses are standard deviations where appropriate. Numbers rounded to nearest whole number. For race: the category Black includes blacks and mulattos and the category Other includes indigenous, Asian and all other. Information on race is not available for Mexico or Argentina (Palloni & Guend, 2005).

support systems, labor market status, and retirement planning. The sample we used in this paper included 12,527 target respondents (no spouses) aged 60 and above who were interviewed in the 2000 wave (Health and Retirement Study, 2002).⁷

Analysis

For self-reported health we use individuals within countries as units of observations and the logit of the probability of self-reporting in bad health as a dependent variable. As predictors we use dummy variables to represent cities, age groups and sex (see definitions in Table 1). Patterns for functional limitations can be studied better with the same tools deployed for the study of self-reported health. We focus on a logit transformation of the individual probabilities of reporting at least one ADL (IADL) and dummy variables for sex, age categories and countries as predictors. To examine the effects of education and income we use logit models while controlling for age, sex and country of origin.

⁷ For more information on the HRS study and sample see the electronic version of papers by Servais (2004) at <http://hrsonline.isr.umich.edu/docs/dmgt/OverviewofHRSPublicData.pdf> or Hauser and Willis (2005) at <http://hrsonline.isr.umich.edu/papers/background/PDR30suppHAUSER.pdf>.

Table 2 Health and other attributes of sample.

Condition/ variable	Overall (<i>n</i> = 10,902)	Argentina (<i>n</i> = 1,043)	Barbados (<i>n</i> = 1,808)	Brazil (<i>n</i> = 2,143)	Chile (<i>n</i> = 1,306)	Cuba (<i>n</i> = 1,905)	Mexico (<i>n</i> = 1,247)	Uruguay (<i>n</i> = 1,450)
Diabetes	17%	13%	22%	18%	14%	15%	22%	13%
Medicine	68%	64%	78%	64%	70%	60%	82%	52%
Insulin	12%	10%	15%	14%	6%	15%	9%	11%
Cancer	4%	5%	4%	4%	5%	3%	2%	6%
Respiratory	10%	8%	4%	13%	13%	13%	10%	9%
Heart	21%	20%	12%	21%	34%	24%	10%	23%
Stroke	7%	5%	6%	8%	7%	10%	5%	4%
Arthritis	42%	53%	47%	33%	32%	58%	25%	47%
Obesity	24%	–	24%	20%	30%	14%	30%	34%
Poor health	11%	5%	5%	9%	21%	13%	20%	7%
ADL	20%	19%	14%	24%	26%	21%	19%	17%
IADL	29%	29%	23%	40%	32%	28%	29%	17%
Height (cm)	158 (10)	–	163 (10)	157 (9)	155 (10)	158(10)	154 (9)	160 (9)
Knee height (cm)	50 (5)	–	53 (5)	50 (3)	48 (3.3)	50 (5)	49 (4)	48 (6)
Weight (kg)	67 (16)	–	72 (20)	64 (13)	67 (14)	61 (14)	66 (12)	72 (15)
BMI (w/h2)	27 (6)	–	27 (8)	26 (5)	28 (5)	25 (5)	28 (5)	28 (7)

Source: SABE data. Numbers rounded to nearest whole number. Poor health 1 = Poor, 0 = All other. ADL = at least 1 ADL. IADL = at least 1 IADL. No height and weight measurements were taken in Argentina.

Results

Self Reported Health

We begin with an assessment of self-reported health status. There is good evidence suggesting that self-reported health is an indicator of general health with good construct validity (Smith, 1994; Manton, Stasllard, & Cordel, 1997; Wallace, 1995; Soldo & Hill, 1995), and is a respectably powerful predictor of mortality risks (Idler & Benyamini, 1997; Idler & Kasl, 1991), disability (Idler & Kasl, 1995) and morbidity (Schechter, Beatty, & Willis, 1998; Beckett, Weinstein, Goldman, & Yu-Hsuan, 2000), though these properties vary somewhat with national or cultural contexts (Idler & Benyamini, 1997). Less is known, however, about whether and to what degree self-reported health status is contaminated by cultural idiosyncrasies, heterogeneous conceptualization of disease and ill health, and differential assessment of gradations of ill health. Even less is known about the impact of these distortions on the validity of direct cross-cultural comparisons of self-reports (Sen, 2002).

Robustness of comparisons: necessary conditions The questionnaires used in SABE were strictly comparable. In particular, the questions eliciting self-reported health status were the same across countries. By the same token, with the exception of Barbados, all countries represented in the study belong to closely related cultural pools which makes it more likely that, if any, the influence of cultural idiosyncrasies will be attenuated. However, these conditions offer only weak assurances for the validity of the cross-country comparability we are about to undertake. In order to verify its robustness we first assess the degree of concordance of self-reported health and self-reported conditions. In particular, we demonstrate that there is a moderate degree of consistency between self-reported health, ADL, IADL and chronic conditions. Figure 1a displays the proportion of respondents with at least one ADL and with at least one IADL by self-reported health status. Figure 1b displays the mean number of self-reported chronic conditions by self-reported health status.




Figure 1 (a) Relation self-report and ADL/IADL, (b) Relation self-report and # chronic conditions, (c) Relation self-report and diabetes.

Finally, figure 1c shows the relation between self-reported health status and self-reported diabetes. The relations portrayed in these graphs are by no means perfect but reveal a high degree of concordance which is satisfying for our purposes.⁸ This is overwhelmingly confirmed by simple descriptive models relating selected health outcomes to self-reported health (Palloni & McEniry, 2004). Regardless of country, the proportion of individuals declaring themselves in bad health is the first or second best predictor of the proportion with at least one ADL, at least one IADL, the mean number of chronic conditions and self-reported diabetes when age, sex and country are controlled for. Exactly the same results hold for the US population in the HRS sample.

This evidence suggests that in each country the proportion self-reporting in bad health reflects underlying medical conditions and functional limitations identified by respondents. Although it is certainly not sufficient for an unbiased cross-national comparison, it signals that we have a relatively robust basis for interpretation of intercountry heterogeneity of self-reports.

⁸ To simplify analyses we focus on a single indicator for the presence of ADL and IADL, namely, whether or not individuals declare at least one of them. We could have used the entire frequency distribution and worked instead with the “number of ADL” or the “number of IADL.” But this complicates the analyses unnecessarily since these are discrete, bounded variables and their distribution can only be mimicked by a handful of discrete distributions. Treating them as categories leads to unwieldy results. Finally, because the number of possible ADL (6) and IADL(6) is relatively few, the proportion of individuals declaring 0 turns out to be an excellent predictor of the shape of the entire distribution. Inferences drawn with the simplified indicator chosen here do not change if the dependent variables are fine-tuned (Palloni & McEniry, 2004). The same applies for self-reported health status.

Heterogeneity of self-reported health Figure 2a displays the proportions reporting their health as “bad” (“mala”) by age groups for all seven cities separately by gender. As a contrast we have included quantities for elderly aged 60 and over who are participants in the Health and Retirement Survey (HRS). The *first* feature of the graph is the massive intercountry heterogeneity which completely overwhelms the effects of gender and age. The cities with the highest proportions of individuals in bad health are found in Santiago (21%), Mexico City (20%) and Havana (13%) whereas those with the lowest are Buenos Aires, Bridgetown, and Montevideo (5 to 7%). The latter three cities are located in countries that, perhaps not coincidentally, are those which until the beginning of the XX1st century enjoyed highest standards of living (as measured by GNP per capita). They are also those with the most modern demographic regime, with near replacement fertility and life expectancies at birth exceeding 75 years.

The *second* feature is the age and sex patterns of self-reports. By and large we observe increasing proportions in bad health with age. The only exception to this regularity is in Mexico City, where the age pattern is flat. The sharpest increase always occurs after age 70 and is particularly pronounced except perhaps in Chile and Mexico. Females do worst than males everywhere, a recurrent finding with data of this type. The Latin American and Caribbean region is no different than others in this respect.


A *third* feature is that elderly living in the cities with the best standing (Buenos Aires, Bridgetown and Montevideo) are equally or less prone than those in the US to report their health as bad, while elderly living in Santiago, Havana and Mexico City are considerably more likely to do so. Elderly in the US are in an intermediate position between these two extremes.

Some of these visual regularities are confirmed by straightforward multivariate analyses of the proportion self-reporting in bad health. The results are presented in Table 3 (Panel a) showing estimates from the most complete model. The results from Model II (Panel b, country effects are constrained to be equal to each other for all countries except the residual category represented by Uruguay) indicate that there is substantial intercountry heterogeneity. Also Model I (Panel a, unconstrained country effects) adds significantly to the fit of Model II shows that the inclusion of the contrast Uruguay versus all other countries adds significantly to the fit of the model (log likelihood ratio test between models leads to a chi-square statistic equal to 116.1 with 6 degrees of freedom). Gender differentials exert a respectable influence whereas those of age are more irregular. In model I only the effects of age groups 75–79 and 85+ are significantly different from 0 but they are not significantly different from each other.⁹ This indicates that there is no firm basis to infer the existence of an age gradient in the proportion of individuals reporting in bad health. Not unlike the pattern found in HRS, where age effects are significantly different from zero but not from each other.

Model III (Panel c) seeks to verify whether or not there are relevant differences between the US and the seven cities in Latin America and the Caribbean. In order to do this we treat the data from the seven cities in Latin America and the Caribbean as if they belonged to the same population and contrast it with the data from HRS. We first estimate a model including age and sex as variables to establish an average age–sex pattern. We then introduce a dummy variable to distinguish the HRS data from the rest. The results suggest that visual impressions can be misleading as elderly people in the US report themselves in

⁹ This statement is established by estimating a model where the age effects are constrained to be the same.

Figure 2 (a) Proportion bad health by age/sex, (b) Proportion ADL by age/sex, (c) Proportion IADL by age/sex.



somewhat better health status than the average of the seven cities in Latin America and the Caribbean. The odds of reporting in bad health among elderly in the US is only about 0.73 as large as among the pooled sample and the implied effect on the log odds of self-reporting in bad health (-0.31) is significantly different from 0 ($t = -6.82$; $p > 0.000$).¹⁰

¹⁰ Simple analyses of variance (Palloni & McEniry, 2004) reveal that the residual variance explained by country heterogeneity is significant whereas the residual variance explained by age and sex is not.

Table 3 Age, gender and country effects on poor self reported health.¹

	Panel A		Panel B		Panel C	
	Effect	SE	Effect	SE	Effect	SE
Constant	-3.00	(0.13)***	-2.95	(0.13)***	-2.41	(0.06)***
Female	0.35	(0.07)***	0.37	(0.07)***	0.33	(0.05)***
65–69 years	0.07	(0.09)	0.04	(0.09)	0.05	(0.06)
70–74 years	0.03	(0.10)	-0.06	(0.10)	-0.04	(0.07)
75–79 years	0.29	(0.10)**	0.18	(0.10)	0.30	(0.07)***
80–84 years	0.20	(0.12)	0.10	(0.11)	0.25	(0.08)**
85+	0.34	(0.13)**	0.24	(0.12)*	0.42	(0.09)***
Argentina	-0.25	(0.18)				
Barbados	-0.22	(0.15)				
Brazil	0.36	(0.13)**				
Chile	1.35	(0.13)***				
Cuba	0.77	(0.13)***				
Mexico	1.28	(0.13)***				
HRS					-0.31	(0.04)***
SABE			0.65	(0.11)***		
<i>N</i>	10,679		10,679		23,200	
Log likelihood	-3,533		-3,691		-7,290	
LR chi square	399		82		155	
Degrees of freedom	12		7		7	

Numbers in parentheses are standard errors. Significance: * $p < 0.05$ ** $p < 0.01$ *** $p < 0.001$.

¹ In this article we defined bad health in terms of elderly who responded “poor” to an assessment of their general health: “Would you say that your health is excellent, very good, good, fair or poor?”

Functional Limitations

Self reported limitations in Activities of Daily Living (ADL's) or Instrumental Activities of Daily Living (IADL's) are a mainstay of population-based information on disability. They are arguably better gauges to assess the extent of physical impairment in population-based studies and are widely used in national surveys such as the HRS, NHANES, NHIS and LSOA as well as in a number of surveys in countries other than the US. ADL's reflects impairments associated with underlying conditions that induce physiological limitations and deterioration and provide a useful benchmark to calibrate demand for care, assistance, and support. IADL's are less tied to morbidity per se as they are sensitive to more generalized impairments and limitations in unassisted and independent living. ADL's are good probes of physical functioning, particularly lower body functionality (Smith, Branch, & Scherr, 1990), and reflect impairment created by chronic conditions as well as cognitive and affective functioning (Stump, Clark, Johnson, & Wolinsky, 1977; Wray, Herzog, & Park, 1996; Wary & Lynch, 1998). As before, we focus only on the proportion of elderly with at least one ADL, or at least one IADL, and examine patterns of relations with the aid of straightforward and parsimonious models.¹¹

The age patterns of the proportions with at least one ADL and at least one IADL are displayed in figure 2b and c. There are strong age gradients, important gender differences, but virtually

¹¹ See Footnote 5.

Table 4 Age, gender and country effects on reporting at least one ADL.

	Panel A		Panel B		Panel C	
	Effect	SE	Effect	SE	Effect	SE
Constant	-2.37	(0.10)***	-2.37	(0.10)***	-2.17	(0.05)***
Female	0.45	(0.05)***	0.45	(0.05)***	0.44	(0.04)***
65–69 years	0.16	(0.08)	0.15	(0.08)	0.11	(0.06)
70–74 years	0.28	(0.09)***	0.26	(0.09)**	0.30	(0.06)***
75–79 years	0.76	(0.08)***	0.75	(0.08)***	0.63	(0.06)***
80–84 years	1.15	(0.09)***	1.14	(0.08)***	1.04	(0.06)***
85+	1.74	(0.09)***	1.71	(0.09)***	1.59	(0.06)***
Argentina	0.14	(0.11)				
Barbados	-0.36	(0.10)***				
Brazil	0.27	(0.09)**				
Chile	0.48	(0.10)***				
Cuba	0.13	(0.09)				
Mexico	0.23	(0.10)*				
HRS					0.10	(0.03)**
SABE			0.15	(.08)		
N	10,824		10,824		21,322	
Log likelihood	-5,041		-5,086		-10,272	
LR chi square (12 <i>df</i>)	706		617		1,082	
Degress of freedom	12		7		7	

Numbers in parentheses are standard errors. Significance: * $p < 0.05$ ** $p < 0.01$ *** $p < 0.001$.

no intercity heterogeneity. The HRS sample stands out: females at older ages in this sample experience much lower proportions with at least one IADL than other SABE females with the exception of Uruguay. Differences noted between US and SABE males tend to be smaller.

Table 4 (panels a,b and c) displays results for ADL and Table 5 (panels a, b and c) does so for IADL. For ADL age and sex patterns are very salient, more so than intercountry heterogeneity.¹² In this case intercountry differences are confined to the contrast between one country with low levels of prevalence of ADL (Barbados) and three with relatively higher levels (Brazil, Chile and Mexico). But, as revealed by Model II (panel b, constrained model), intercountry heterogeneity is trivial as the effects associated with all SABE countries (except Uruguay which is the residual category) are not significantly different from 0. Also note that Model I does not add significantly to the fit (relative to Model II). Instead for IADL, intercountry differences are powerful and generalized. A curious feature is that the city with one of the lowest proportion of elderly reporting themselves in poor health has one of the highest proportions with at least one ADL or IADL (Montevideo).¹³

The contrasts between ADL and IADL patterns in SABE cities and HRS (see Panels c in Tables 4 and 5) are quite strong. An individual in the HRS population is about 1.11 times as likely to experience at least one ADL: as an individual in the pooled SABE sample. In fact, the estimated effect on the log odds is equal to 0.10 and is significantly different form zero ($t = 2.93, p > 0.01$). By contrast, the HRS population is less likely (log odds equal to .54) to

¹² Analyses of variance (Palloni & McEniry, 2004) suggest that the fraction of total variance explained by country variability is statistically insignificant.

¹³ Montevideo is also the only city in the SABE sample where institutionalization of the elderly is more than trivial. The peculiar relation between self-reported health and ADL and IADL in Montevideo might be a result of heavy selection among elderly who remain independent instead of becoming institutionalized.

Table 5 Age, gender and country effects on reporting at least one IADL.

	Panel A		Panel B		Panel C	
	<i>Effect</i>	<i>SE</i>	<i>Effect</i>	<i>SD</i>	<i>Effect</i>	<i>SE</i>
Constant	-3.81	(0.13)***	-2.81	(0.09)***	-2.45	(0.06)***
Female	0.59	(0.06)***	0.55	(0.06)***	0.43	(0.04)***
65–69 years	0.35	(0.10)***	0.30	(0.10)**	0.17	(0.07)**
70–74 years	0.67	(0.10)***	0.62	(0.10)***	0.36	(0.07)***
75–79 years	1.21	(0.10)***	1.23	(0.09)***	0.89	(0.07)***
80–84 years	1.72	(0.10)***	1.74	(0.10)***	1.38	(0.07)***
85+	2.60	(0.10)***	2.59	(0.10)***	2.05	(0.07)***
Argentina	0.81	(0.14)***				
Barbados	0.66	(0.12)***				
Brazil	1.49	(0.11)***				
Chile	1.11	(0.13)***				
Cuba	0.85	(0.12)***				
Mexico	1.11	(0.13)***				
HRS					-0.61	(0.04)***
N	10,798		10,798		23,303	
Log likelihood	-4453		-4576		-8768	
LR chi square	1413		1168		1700	
Degrees of freedom	12		6		7	

Numbers in parentheses are standard errors. Significance: * $p < 0.05$ ** $p < 0.01$ *** $p < 0.001$. Panel C is a pooled HRS-SABE sample. For this analysis, we used only those SABE IADLs that were strictly comparable with HRS IADLs: preparing meals, managing money, buying food or clothing, using the phone, and taking medicine.

declare at least one IADL as they are in the pooled sample. The associated effects on the log odds (-0.61) is again significantly different from 0 (-0.61; $t = -15.48$; $p > 0.000$). This is an interesting pattern which could result from heavier mortality selection among elderly with compromising morbid conditions in the SABE cities.

Chronic Conditions: The Salience of Diabetes


Figure 3 displays the mean number of chronic conditions by age and gender shows that, as was the case for ADL and IADL, the age patterns slope upwards and females exhibit a more unfavorable profile than males. By the same token, a simple regression analysis reveals that intercountry heterogeneity is quite low and that the strongest effects are those of age and sex. A comparison with HRS shows that the elderly population of HRS exhibits a higher average number of chronic conditions than *any* of the countries in the SABE sample (except perhaps Cuban females). This again is a pattern that could be expected under heavier mortality selection in SABE countries.¹⁴

Of all chronic conditions highlighted in Table 2 and those included in the mean number of chronic conditions in figure 3, arthritis, heart disease, obesity and diabetes are the most salient.¹⁵ Of these, the latter three are of particular interest to us. First, other research reports

¹⁴ See Appendix for definition of chronic conditions.

¹⁵ In this paper we reserve the term diabetes to refer to a mixture of diabetes 1 and diabetes mellitus or type 2. However, for the most part those individuals self-reporting diabetes are afflicted by diabetes type 2.

Figure 3 Number of chronic conditions by age/sex.



that developing countries, particularly Latin America and the Caribbean, are in the midst of a diabetes (and obesity) epidemic, in part the result of a unfavorable shift toward “Western” diet, rich in saturated fats, simple carbohydrates and sugar and a marked trend toward sedentarism (Popkin, 1993; Albala et al., 2000). But never before has this been documented on a large scale for countries of the region and for elderly populations. Second, diabetes and coronary heart disease have been linked to unfavorable early childhood conditions that express themselves either in unfavorable nutritional status or as results of contraction of infectious diseases (Barker, 1998; Kuh & Ben-Shlomo, 2004). The case of diabetes is of special interest for it, but not heart disease, appears to be strongly related to indicators of early childhood malnutrition (Barker, 1998; Hales & Barker, 1992; Hales et al., 1991; Lithell et al., 1996; Palloni et al., 2004).


Figure 4 displays the proportion of individuals by age groups who self-report diabetes for all SABE samples and HRS, separately by gender.¹⁶ The pattern by age is very distinct and quite similar across countries, concave downward with a peak around 70–74.¹⁷ Males are less likely to report diabetes than females. For the most part, the population in HRS is as likely as the average individual in the pooled SABE sample to report diabetes. In regression models not shown here we find several regularities. First, within the SABE sample there is important heterogeneity. Cuba, closely followed by Argentina and Uruguay, exhibit the lowest levels of self-reported diabetes (almost 17% at the peak age group for women), whereas Barbados, Mexico and Brazil, experience the highest levels (about 29% at the peak age for women). Second, in the HRS sample prevalence of self-reported diabetes is slightly higher than the weighted average in SABE: the predicted probability at the peak age for women is 0.22. But the difference between this level and the prevalence in the SABE sample is statistically insignificant, as the effect of being in the HRS sample on the log odds of self-reporting diabetes is only 0.10 (with $t = 0.73, p > 0.470$).

The fact that Cuba exhibits very low levels of self-reported diabetes specially among males, is undoubtedly due, in part at least, to the fact that the adoption of a Western life style has simply not been an option in this country and, therefore, the risk factors associated with a new diet and sedentary life styles are simply absent. But to explain the very high

¹⁶ Self-reports of diabetes are not perfect. We know that it underestimates true prevalence. But it is quite accurate as it has very high specificity but lower sensitivity in very different cultural contexts (Palloni, Soldo, & Wong, 2003; Goldman, I-Fen, Weinstein, & Yu-Hsung, 2002).

¹⁷ The declining pattern with age is probably a result of the heavier attrition of diabetics as age increases.

Figure 4 Proportion diabetes by age/sex.



levels in Barbados, Brazil and Mexico one probably needs to explore the role of population composition by early nutritional status and/or the influence of ethnic composition. In fact, Barbados and Brazil have a hefty component of population with African descent whereas Mexico has the highest percentage of indigenous and mestizo population. Whether the fetal origin explanation or the one resting on ethnic-related genetic endowment or a combination of both explain the distinctive patterns in these countries must remain conjectures until we are able to test directly the influence exerted by each factor.¹⁸

Health Inequalities

The most important conjecture we put forward at the outset is that the evolution of mortality in countries of the region may have important implications for the health status of those entering older age brackets now and in the next 20 to 30 years. In particular, if deleterious early childhood conditions do indeed affect health status during adulthood, we would expect that individuals who belong to more privileged social groups would be less affected by the peculiar evolution of mortality trends early during the century, and should experience a more favorable profile than those in less privileged social positions. Thus, one possible, albeit indirect, test of the conjecture should rely on contrasts of health status by socioeconomic standing. However, comparisons of health status across social groups are far from an ideal test for at least two reasons. First, similar contrasts in health status by social classes can also be the result of other mechanisms, such as better access to health care or more favorable behavior. For example, one would expect inequalities to be more muted in Cuba than elsewhere. This does not mean that the conjecture is incorrect; only that other mechanisms help to conceal its effects. Second, since average frailty of those in better socioeconomic standing is likely to be higher than those in lower socioeconomic positions, health differentials at older ages will be considerably attenuated and will make less visible the contrasts associated with early experiences. Thus, for example, social classes who experienced higher mortality during childhood may have lost a larger fraction of individuals who would be in worse health had they survived, as they did among better off social classes. This mechanism alone can produce weaker contrasts and be more powerful in countries where social contrasts in morbidity and mortality were stronger early in the life of

¹⁸ See Palloni and McEniry, 2004. Since most of the Barbados population is of African descent, a similar test cannot be applied there.

the cohorts of elderly. These factors with offsetting effects may blur distinctions and make it difficult to identify the presence of any particular mechanisms.

The SABE data set includes a number of variables that can be used to assess socioeconomic status. We will concentrate in two that are least controversial, levels of attained education and family income. Neither is an ideal measure of social class but both are strongly associated with mortality and morbidity in other social contexts and, in all likelihood, reflect a number of conditions with powerful effects on health status. Figure 5a and b graphically display the aggregate association between education (income) and selected health outcomes. These figures plot the proportion experiencing a particular outcome by quintiles of education (income). In both cases the relations seem to be in the right direction, and the slopes of gradients are in the expected direction. There is some intercountry heterogeneity, and Cuba exhibits the lowest levels of inequality across quintiles of income and education categories. But the differences are trifling. Inequality by income or education is more pronounced for self-reported health, ADL and IADL in that order and least pronounced for obesity. There are some observable social inequalities for diabetes but probably they betray much higher levels concealed by differential mortality of diabetics across social classes.

A summary of the effects for the SABE sample appears in Table 6 (panel a). The effects of both education and income are pervasive, powerful and statistically significant for almost all health outcomes except those of income on diabetes and obesity. When we contrast these inequalities with those in HRS (Table 7, panel b) we find a striking pattern: HRS inequalities by both education and income are larger in virtually all health outcomes than those found in SABE countries.




Figure 5 (a) Association between education & health outcomes, (b) Association between income & health outcomes.

Table 6 (panel a): Education and income effects on morbidity controlling for age, gender and country (SABE only).

	<i>n</i>	Effect	SE	Log likelihood	Chi-square with (<i>df</i>)
Education					
Self report					
Secondary	9,584	-0.55	(0.10)***	-2,981	369 (<i>df</i> = 14)
Higher education		-0.89	(0.16)***		
ADL					
Secondary	9,732	-0.46	(0.08)***	-4,365	661 (<i>df</i> = 14)
Higher education		-0.60	(0.11)***		
IADL					
Secondary	9,688	-0.36	(0.07)***	-4,838	1484 (<i>df</i> = 14)
Higher education		-0.59	(0.20)***		
Diabetes					
Secondary	9,751	-0.17	(0.07)*	-4,301	164 (<i>df</i> = 14)
Higher education		-0.52	(0.11)***		
Obesity					
Secondary	7,862	-0.18	(0.07)**	-4,002	677 (<i>df</i> = 13)
Higher education		-0.41	(0.10)***		
Income					
Self Report					
Middle	8,864	-0.19	(0.08)*	-2,893	382 (<i>df</i> = 14)
Richest		-0.78	(0.12)***		
ADL					
Middle	9,002	-0.05	(0.07)	-4,193	625 (<i>df</i> = 14)
Richest		-0.34	(0.09)***		
IADL					
Middle	8,972	-0.18	(0.06)**	-4,555	1604 (<i>df</i> = 14)
Richest		-0.63	(0.08)***		
Diabetes					
Middle	9,009	-0.05	(0.07)	-3,950	145 (<i>df</i> = 14)
Richest		-0.13	(0.09)		
Obesity					
Middle	7,331	0.11	(0.07)	-3,712	579 (<i>df</i> = 13)
Richest		0.01	(0.09)		

Significance: * $p < 0.05$ ** $p < 0.01$ *** $p < 0.001$.

Are these effects important? Although statistically significant effects are worth studying, they not always translate into important influences. To illustrate the magnitude of the actual effects we calculate predicted probabilities of self-reported diabetes for individuals with average attributes except in education and income. In the SABE cities the average probability of diabetes increases from about 0.14 to 0.18 when education levels changes from high school and above to primary (complete or incomplete). In HRS the shift is from 0.16 to 0.23. Thus a change in educational level of this sort doubles the probabilities of diabetes in both the US and Latin American context. For income the contrasts are slightly more attenuated in SABE: the average chance of diabetes for those in the lowest quintile of the income distribution is about 1.20 as high as the chance for those in the highest quintile (0.18 against 0.15). In HRS the ratio of one to the other is slightly higher than 2 (0.23 to 0.11).

Table 7 (panel b): Education and income effects on morbidity controlling for age and gender (HRS only).

	<i>n</i>	Effect	SE	Log likelihood	Chi-square (with <i>df</i>)
Education					
Self report					
Secondary	12,500	-1.18	(0.09)***	-3,424	368 (<i>df</i> = 8)
Higher education		-1.95	(0.11)***		
ADL					
Secondary	10,480	-0.80	(0.08)***	-5,104	613 (<i>df</i> = 8)
Higher education		-1.09	(0.09)***		
IADL					
Secondary	12,484	-0.78	(0.09)***	-4,076	495 (<i>df</i> = 8)
Higher education		-1.24	(0.10)***		
Diabetes					
Secondary	12,504	-0.32	(0.09)	-5,520	130 (<i>df</i> = 8)
Higher education		-0.76	(0.09)***		
Obesity					
Secondary	12,358	-0.34	(0.09)***	-6,249	453 (<i>df</i> = 8)
Higher education		-0.73	(0.09)***		
Income					
Self report					
Middle	12,473	-1.11	(0.07)***	-3,339	499 (<i>df</i> = 8)
Richest		-2.46	(0.16)***		
ADL					
Middle	10,461	-0.76	(0.06)***	-5,015	771 (<i>df</i> = 8)
Richest		-1.38	(0.09)***		
IADL					
Middle	12,458	-0.93	(0.07)***	-4,002	624 (<i>df</i> = 8)
Richest		-1.43	(0.11)***		
Diabetes					
Middle	12,477	-0.55	(0.06)***	-5,475	204 (<i>df</i> = 8)
Richest		-1.03	(0.08)***		
Obesity					
Middle	12,331	-0.33	(0.06)***	-6,232	449 (<i>df</i> = 8)
Richest		-0.68	(0.07)***		

Significance: * $p < 0.05$ ** $p < 0.01$ *** $p < 0.001$.

Discussion

This review of preliminary findings from SABE reveals well-known patterns and confirms routine expectations and, simultaneously, presents us with puzzling regularities that do not agree well neither with a priori expectations nor with conjectures. At the outset we posed three questions. The first question had to do with the characteristic health profile of the elderly in the region. We now know a few things we did not know before. Self reported health status shows large intercountry variability and more muted heterogeneity due to gender and age. Women and the very old are more likely to declare themselves in bad health. On average, countries in the region display patterns by age and sex that are very similar to those found elsewhere. The proportion with at least one ADL or IADL is strongly related to age and gender and displays remarkable intercountry invariance. Self reported health, on the one hand, and ADL and IADL on the other are moderately related to each other. The mean number of self-reported chronic conditions increases with age and is higher for females than it is for males. Of all the chronic

conditions the one that stands for its high prevalence is (self reported) diabetes. Females are specially affected by this condition. Obesity, a risk factor of diabetes, is also high among elderly in the region, particularly among females. There is large intercountry variance in both diabetes and obesity. In some countries (such as Argentina) who are highly modernized and westernized, the prevalence of diabetes is fairly low and in others, such as Mexico and Barbados, with high percentages of population of African descent or mestizo, the prevalence is very high.

The second and third questions had to do with the relative standing of health status in the region relative to other countries. In this paper we chose the US as a benchmark not because it is uniquely suited to be so, but because the data were available to us. We find that the health profile of elderly in the Latin America and Caribbean is better in some respects and worse in others. Patterns of self-reported health are comparable whereas the prevalence of at least one IADL tends to be lower in the US than in SABE. Similarly, the US population exhibits a much larger mean number of chronic conditions at all ages and particularly among males. The most interesting feature is that, on the whole, SABE countries, on average, display levels of self-reported diabetes (and obesity) that are as high if not higher than those found in the US.

In the end, do these findings lead to straightforward policy implication? The answer is a resounding yes. Of all the features we discovered, the high prevalence of diabetes and obesity is perhaps the most worrisome. It is known that the health costs implications of diabetes are staggering even if the disease presents itself with a normal, expected profile of associated comorbidities. If lack of compliance with treatment or the mixture of diseases to which the population is exposed complicates the clinical status of individual with diabetes, the costs could escalate even more. Regardless of the root causes, a continued increase in the prevalence of diabetes will pose severe constraints on the health system in these countries. What seem most obvious is that educational campaigns are cost-effective means to both reduce incidence and keep the disease negative spillovers at bay. In the next several decades major health policy initiatives for the elderly will have to include prevention of obesity and diabetes.

Acknowledgments This paper is only possible thanks to the collaboration of the principal investigators of the SABE study, Cecilia Albala, Anselm Hennis, Roberto Ham, Maria Lucia Lebrao, Esther de Leon, Edith Pantelides, and Omar Pratts. We are thankful to Dr. Guido Pinto for many discussions and for extensive work on the data set and to Dr. Martha Pelaez, from the Pan American Health Organization, without whose initiative the project would not have been possible. The research for this paper was supported by NIA grants R01 AG16209 and R03 AG15673 to Palloni. Both authors work in the Center for Demography and Ecology supported by core grant P30 HD05876, and in the Center Core supported by core grant P30 AG17266.

Appendix

ADL and IADL

1.1. ADL's:

- Walking across the room
- Dressing
- Bathing
- Eating
- Getting in and out of bed
- Using bathroom

1.2 . IADL's

- Preparing meals
- Managing money
- Difficulty with getting to places (only in SABE)
- Buying food or clothing
- Using the phone (in SABE asked of those who had a phone)
- Doing heavy housework (only in SABE)
- Doing light housework (only in SABE)
- Taking medicines

For multivariate analyses, we used only those SABE IADLs that were strictly comparable with HRS: preparing meals, managing money, buying food or clothing, using the phone, and taking medicine.

Chronic Conditions

- Arthritis
- Cancer
- Diabetes
- Respiratory Illness
- Heart Disease
- Stroke

Targets, Spouses and Proxies

In three countries (Argentina, Chile, and Uruguay) only one individual per household was interviewed. In two countries, Brazil and Mexico, interviewers proceeded to interview all individuals 60 and older found in selected household. In virtually all these cases, the additional interviews corresponded to spouses (one per household). In Cuba interviewers selected a target individual and a spouse.

In our analyses we include all individuals interviewed. This has the advantage of maximizing observation at the expenses of introducing dependence of observations in the countries where more than one individual per household was interviewed. In order to protect our inferences we repeated some of the analyses using clustering procedures to adjust for lack of independence but since the inferences remain unchanged we have chosen to present results based on the larger samples.

Sampling Weights

Only the sample from Santiago is self-weighted. All others require weights to expand the sample population to the city population. Since in two countries no sample weights have been calculated we chose to ignore them in all the others. However, to ensure that none of our conclusions was sensitive to this choice, we proceeded to re-estimate models using sampling weights for those countries that had them available. None of the inferences changed, and it is highly unlikely that they will even in the countries where there were no weights available yet.

References

- Albala, C., Kain, J., Burrows, R., & Diaz, E. (2000). *Obesidad: Un desafío pendiente*. Santiago, Chile: Editorial Universitaria, Universidad de Chile.
- Albala, C., Lebrao, M. L., Leon Diaz, E. M., Ham-Chande, R., Hennis, A. J., Palloni, A., et al. (2005). The health, well-being and aging (SABE) study: Sample method and profile of the studied population. *Pan American Journal of Public Health*, 17(5/6), 307–322.
- Barker, D. J. P. (1998). *Mothers, babies and health in later life (2nd Edition)*. Edinburgh: Churchill Livingstone.
- Barrientos, A. (1997). The changing face of pensions in Latin America: Design and prospects of individual capitalization pension plans. *Social Policy & Administration*, 31(4), 336–353 (December).
- Beckett, M., Weinstein, M., Goldman, N., & Yu-Hsung, L. (2000). Do health interview surveys yield reliable data on chronic illness among older respondents? *American Journal of Epidemiology*, 151(3), 315–323.
- Cynader, Max S. (1994). Mechanisms of brain development and their role in health and well-being. *Daedalus*, 123(4), 155–165.
- Devos, S. (1990). Extended family living among older people in six Latin American countries. *Journal of Gerontology*, 45(3), S87–S94.
- Devos, S., & Palloni, A. (2002). *Living arrangements of elderly people around the world*. University of Wisconsin-Madison: Center for Demography & Ecology.
- Elo, I. T., & Preston, S. H. (1992). Effects of early-life conditions on adult mortality: A review. *Population Index*, 58(2), 186–212.
- Frenk, J., Frejka, T., Bobadilla, J. L., Stern, C., & Sepulveda, J. (1991). Elements for a theory of the health transition. *Health Transition Review*, 1(1), 21–38.
- Goldman, N., I-Fen, L., Weinstein, M., & Yu-Hsung, L. (2002). *Evaluating the quality of self-reports on hypertension and diabetes*. Princeton University: Office of Population Research.
- Hales, C. N., & Barker, D. J. P. (1992). Type 2 (non-insulin-dependent) diabetes mellitus: The thrifty phenotype hypothesis. *Diabetologia*, 35, 595–601.
- Hales, C. N., Barker, D. J. P., Clark, P. M. S., Cox, L. J., Fall, C., & et al. (1991). Fetal and infant growth and impaired glucose tolerance at age 64. *British Medical Journal*, 303, 1019–1022.
- Hauser, R. M., & Willis, R. J. (2005). http://hrsonline.isr.umich.edu/papers/background/PDR30supp_HAUUSER.pdf. This is an electronic version of an article published in *Aging, Health, and Public Policy: Demographic and Economic Perspectives, a supplement to Population and Development Review Volume 30*. New York: Population Council, 2005.
- Health and Retirement Study, HRS Core (Final) (v 1.0) public use dataset (2002). Produced and distributed by the University of Michigan with funding from the National Institute of Aging, (U01 AGO 9740), Ann Arbor, Michigan.
- Hertzman, C. (1994). The lifelong impact of childhood experiences: A population health perspective. *Daedalus*.
- Idler, E. L., & Benyamini, Y. (1997). Self-rated health and mortality: A review of twenty-seven community studies. *Journal of Health and Social Behavior*, 38, 21–37.
- Idler, E. L., & Kasl, S. (1991). Health perceptions and survival: Do global evaluations of health status really predict mortality? *Journal of Gerontology*, 46(2), S55–S65.
- Idler, E. L., & Kasl, S. V. (1995). Self-ratings of health: Do they also predict change in functional ability? *Journal of Gerontology*, 50B(6), S344–S353.
- Kinsella, K., & Velkoff, V. (2001). *An Aging World*. US Government Printing Office, Washington, District of Columbia: US Bureau of the Census.
- Klinsberg, B. (2000). *America Latina: Una region en riesgo, pobreza, inequidad e institucionalidad social*. Inter American Development Bank.
- Kuh, D., & Ben-Shlomo, Y. (eds). (2004). *A life course approach to chronic disease epidemiology*. Oxford: Oxford University Press.
- Lithell, H. O., McKeigue, P. M., Berglund, L., Mohsen, R., Lithell, U. B., & Leon, D. A. (1996). Relation of size at birth to non-insulin dependent diabetes and insulin concentrations in men aged 50–60 years. *British Medical Journal*, 312, 406–410.
- Manton, K. G., Stallard, E., & Cordel, R. (1997). Changes in age dependence of mortality and disability: Cohort and other determinants. *Demography*, 34(1), 135–157.
- Mesa-Lago, C. (1994). *Changing social security in Latin America: Toward alleviating the costs of economic reform*. Boulder and London: Lynne Rienner.
- Oman, A. R. (1982). Epidemiologic transition. In *International Encyclopedia of Population* (pp. 172–83). New York: Free Press.
- Palloni, A. (2001). Living arrangements of older persons. *United Nations Population Bulletin* 42/43.

- Palloni, A., & Guend, H. (2005). Stature prediction equations for elderly Hispanics by gender and ethnic background developed from SABE data. *Journal of Gerontology: Medical Sciences*, 60(6), 804–810.
- Palloni, A., & McEniry, M. (2004). Health status of elderly people in Latin America. Madison, Wisconsin: Center for Demography and Ecology, University of Wisconsin. Working paper.
- Palloni, A., & Pelaez, M. (2002). *Survey of health and well-being of elders*. Washington, District of Columbia: Pan American Health Organization. Final Report.
- Palloni, A., McEniry, M., Guend, H., Davila, A. L., Garcia, A., Mattei, H., & Sanchez, M. (2004). *Health among Puerto Ricans: Analysis of a new data set*. Center for Demography and Ecology. University of Wisconsin-Madison.
- Palloni, A., Pinto, G., & Pelaez, M. (2002). Demographic and health conditions of ageing in Latin America. *International Journal of Epidemiology*, 31, 762–771.
- Palloni, A., Soldo, B., & Wong, R. (2003). The accuracy of self reported anthropometric measures and self reported diabetes in nationally representative samples of older adults in Mexico. Paper presented at the Population Association of America Minneapolis, Minnesota. May 1–3.
- Palloni, A., & Wyrick, R. (1981). Mortality decline in Latin America: Changes in the structures of causes of deaths, 1950–1975. *Social Biology*, 28(3–4), 187–216.
- Popkin, B. M. (1993). Nutritional patterns and transition. *Population Development Review*, 19, 138–157.
- Preston, S. H. (1976). *Mortality patterns in national populations with special reference to recorded causes of death*. New York: Academic.
- Ruggles, S. (1996). Living arrangements of the elderly in America. In Tamara K. Hareven (ed), *Ageing and generational relations over the life course: A historical and cross-cultural perspective* (pp. 254–271). New York: Walter de Gruyter.
- SABE. Salud y Bienestar en el Adulto Mayor, SABE, version No 1, restricted circulation data set (2003). Produced and distributed by the Pan American Health Organization (PAHO) and the Center for Demography and Health of Aging (CDHA) with the support of the National Institute of Aging, R03 AG15673.
- Schaffer, R. H. (2000). The early experience assumption: Past, present, and future. *International Journal of Behavioral Development*, 24(1), 5–14.
- Schechter, S., Beatty, P., & Willis, G. B. (1998). Asking survey respondents about health status: Judgment and response issues. In N. Schwarz, D. Park, B. Knauper, & S. Sudman, (eds), *Cognition, Aging, and Self-Reports*. Ann Arbor, Michigan: Taylor and Francis.
- Sen, A. (2002). Perception versus observation. *BMJ*, (324), 859–860.
- Servais, M. A. (2004). Overview of HRS public data files for cross-sectional and longitudinal analysis. Survey Research Center, Institute for Social Research, University of Michigan, Ann Arbor.
- Smith, J. (1994). Measuring health and economic status of older adults in developing countries. *Gerontologist*, 34(4), 491–496.
- Smith, L. A., Branch, L. G., & Scherr, P. A. (1990). Short-term variability of measures of physical function in older people. *Journal of American Geriatric Society*, 38, 993–998.
- Soldo, B. J., & Hill, M. (1995). Family structure and transfer measures in the health and retirement study: Background and overview. *Journal of Human Resources, Supplement*, 108–137.
- Stump, T. E., Clark, D. O., Johnson, R. J., & Wolinsky, F. D. (1977). The structure of health status among Hispanic, African American, and white older adults. *Journals of Gerontology*, 52B, 49–60 (Special Issue).
- United Nations (2000). *United Nations demographic yearbook*. United Nations, New York: Department of Social and Economic Affairs, Table 7.
- Vaupel, J., Manton, K., & Stallard, E. (1979). The impact of heterogeneity in individual frailty on the dynamics of mortality. *Demography*, 16(3), 439–454.
- Wallace, R. B. H. A. R. (1995). Overview of the health measures in health and retirement study. *Journal of Human Resources, Supplement*, 84–107.
- Wray, L. A., Herzog, A. R., & Park, D. C. (1996). Physical health, mental health, and function among older adults. Paper presented at the Annual Meetings of the Gerontological Society of America, November, Washington, District of Columbia.
- Wray, L. A., & Lynch, J. W. (1998). The role of cognitive ability in links between disease severity and functional ability in middle-aged adults. Paper presented at the Annual Meetings of the Gerontological Society of America, November, Philadelphia.